

• REVIEW •

# Influencing factors of pancreatic microcirculatory impairment in acute pancreatitis

Zong-Guang Zhou, You-Dai Chen

Zong-Guang Zhou, You-Dai Chen, Department of Hepato-bilio-pancreatic Surgery & Institute of Microcirculation, West China Hospital, Sichuan University, Chengdu 610041, Sichuan Province, China

Correspondence to: Dr. Zong-Guang Zhou, Department of Hepato-bilio-pancreatic Surgery & Institute of Microcirculation, West China Hospital, Sichuan University, Chengdu 610041, Sichuan Province, China. 258836@mail.guoli.com.cn

Telephone: +86-28-5422484

Received 2001-11-02 Accepted 2001-12-05

## Abstract

**Pancreatic microcirculatory disturbance plays an important role in the pathogenesis of acute pancreatitis, and it involves a series of changes including vasoconstriction, ischaemia, increased vascular permeability, impairment of nutritive tissue perfusion, ischaemia/reperfusion, leukocyte adherence, hemorrhological changes and impaired lymphatic drainage. Ischaemia possibly acts as an initiating factor of pancreatic microcirculatory injury in acute pancreatitis, or as an aggravating/continuing mechanism. The end-artery feature of the intralobular arterioles suggests that the pancreatic microcirculation is highly susceptible to ischaemia. Various vasoactive mediators, as bradykinin, platelet activating factor, endothelin and nitric oxide participate in the development of microcirculatory failure.**

Zhou ZG, Chen YD. Influencing factors of pancreatic microcirculatory impairment in acute pancreatitis. *World J Gastroenterol* 2002;8(3):406-412

## INTRODUCTION

Acute pancreatitis remains an important surgical problem with high morbidity and mortality<sup>[1-4]</sup>. It is not merely an injury caused by the activated pancreatic enzymes but also involves pancreatic ischaemia. Evidences in basic and clinical research suggest that disturbance of pancreatic microcirculation plays an important role in its pathophysiological processes<sup>[5-14]</sup>. The specific local microcirculatory changes cannot be prevented merely by adequate fluid therapy. In recent years, studies with modern molecular biological tools have elucidated that many factors are involved in the development of pancreatic microcirculatory disturbance. Whether the disturbance of pancreatic microcirculation is an initiating factor or as a consequence of progressive pancreatitis is still debatable. The pathophysiological changes of pancreatic microcirculatory disturbance in acute pancreatitis are complex, they include local release of acinar enzymes<sup>[15-25]</sup>, vasoactive mediators<sup>[26-39]</sup>, vasoconstriction, increase in vascular permeability, ischaemia<sup>[40-41]</sup>, ischaemia/reperfusion, leukocyte adherence, intravascular coagulation, capillary stasis, etc., resulting in pancreatic oedema, hemoconcentration, and impaired capillary and venous drainage<sup>[42-44]</sup>, consequently leading to hemorrhagic pancreatic necrosis<sup>[45]</sup>.

## ROLES OF ISCHAEMIA IN PANCREATIC MICROCIRCULATORY DISTURBANCE DURING ACUTE PANCREATITIS

### *Ischaemia as an initiating factor*

There is a considerable evidence supporting ischaemia as an initiating

factor of pancreatic microcirculatory injury in acute pancreatitis<sup>[46-48]</sup>. As long ago as 1862, Panum induced hemorrhagic pancreatitis by injection of wax droplets into pancreatic arteries. Later similar changes were noticed by intra-arterial injection of 8-20µm microspheres, irreversibly obstructing terminal arterioles and occluding the capillaries. While the use of larger particles only results in pancreatic oedema, because there are abundant arcade-like anastomoses between the pancreatic interlobular vessels. There is also evidence suggesting that microvascular injection of microspheres may progress to chronic active pancreatitis.

A clinical report revealed at autopsy that atheromatous thrombi embolized from the aorta into the pancreatic arteries were associated with acute pancreatitis in 10 of 12 cases. The incidence of pancreatitis in 182 patients died after cardiac surgery was 16%. There was also evidence for a high susceptibility of the pancreas to ischaemic injury in patients died of shock. A high incidence of acute pancreatitis shown after cardiopulmonary bypass operations seemed to be associated with intraoperative hypoperfusion in the splanchnic area.

By means of intravital microscopy in conjunction with technique of selected cells-labeling, direct impairments of pancreatic microcirculation in the early phase of acute pancreatitis have been observed in the experimental ischaemia induced by controlled haemorrhage or interruption of arterial blood supply to the pancreas<sup>[49]</sup>, suggesting the pancreatic microcirculation being highly susceptible to ischaemia. This is closely related to the microvasculature of pancreatic lobule; there is a single centrally-located intralobular artery as the exclusive vascular supply of each lobule, no anastomosis between the intralobular arteries and their branches exists, indicating the cause of its high susceptibility to ischaemia<sup>[50,51]</sup>.

### *Ischaemia as an aggravating and continuing mechanism*

Temporary complete or partial ischaemia of pancreas would not cause hemorrhagic pancreatic necrosis, the slight histological and functional changes are completely reversible. However, temporary ischaemia has the potential of being transitional from edematous to necrotizing pancreatitis<sup>[52]</sup>. While temporary arterial occlusion alone does not injure the pancreas following induction of edematous pancreatitis by duct ligation<sup>[53]</sup> with hyperstimulation, arterial occlusion for only 15 min can result in parenchymal necrosis, suggesting that ischaemia as an aggravating factor participates in the development of acute pancreatitis.

Impairment of microcirculatory perfusion of pancreas is the consequence of the effect of various local factors, such as vasoconstriction, free radicals, intravascular coagulation, release of vasoactive mediators taking part in the whole course of acute pancreatitis (see below). Recently, ischaemia/reperfusion is considered one of the important causative factors for development of acute pancreatitis after pancreatic transplantation. It has been repeatedly demonstrated that change of pancreatic perfusion is an early event in experimental acute pancreatitis, and microcirculatory impairment in human pancreas also correlate well with the degree of ischaemic injury. These findings support the hypothesis that the microvasculature is the primary target of reperfusional injury after ischaemia.

## CHANGES OF PANCREATIC MICROCIRCULATION IN ACUTE PANCREATITIS

Many indirect methods have been applied to assess the changes of pancreatic microcirculation during acute pancreatitis in previous studies<sup>[54]</sup>. Recently, intravital fluorescence microscopy combined with the technique of separate labeled-cells and computerized image analysis system has been successfully used in the studies of pancreatic microcirculation in acute pancreatitis. Many important phenomena as vascular permeability change, vasoconstriction, capillary blood flow, functional capillary density, leukocyte-endothelium interaction, etc., have been continuously and directly observed during the course of acute pancreatitis. It is now believed that microcirculatory changes are important as well as early feature in the pathophysiology of acute pancreatitis<sup>[55]</sup>.

### Vasoconstriction

The first step in the sequence of microcirculatory events in pancreatitis is the constriction of interlobular vessels, especially in the proximal segments of the interlobular arterioles and venules<sup>[56,57]</sup>. The vasoconstriction occurring in the early phase of acute pancreatitis may cause ischaemia and stasis of the microcirculation<sup>[58]</sup>, which can be prevented by the radical scavengers, superoxide dismutase and N-(2-mercaptopropionyl)glycine in sodium taurocholate-induced pancreatitis, suggesting that vasoconstriction might be induced by free radicals. There is also great support for the concept that solutions injected into the pancreatic duct to induce biliary pancreatitis exert their effect via the interstitial route. Even at a low injection pressure of 40 cmH<sub>2</sub>O, rupture of the ducto-acinar junction is detectable with subsequent fluid extravasation in the interstitial space, where they gain access to the pancreatic microvasculature precipitating vascular spasm. It has been noticed that segmental constriction of pancreatic arteries occurred in bile-induced pancreatitis, and of mesenteric arteries directly exposed to diluted bile. There is also pronounced damage to the pancreatic vessels resulting in haemorrhage, endothelial detachment and thrombosis, as has been shown with the taurocholate, trypsin, and trypsin-digested blood vessels. The vasotoxic effect of these substances was further substantiated by the demonstration that interstitial injection into the omentum precipitates similar changes at the injection site. The finding that stress and shock can convert oedematous to hemorrhagic experimental pancreatitis suggests that catecholamines mediators might participate in the process. Therefore, pancreatic vasoconstriction in acute pancreatitis might be relevant to a variety of factors.

### Changes of permeability

The intravital microscopic findings of immediate leakage of the macromolecular plasma marker (FITC-Dextran 70) from the microvasculature into the interstitial tissue, and the scanning electron microscopic evidence of leakage of the cast material through the capillary membrane in the early phase of acute experimental pancreatitis suggest that presence of increased permeability during the disease process. Further experiments demonstrate that permeability changes precede stasis and stasis precedes leukocyte adherence<sup>[59]</sup>, suggesting that increased vascular permeability and ischaemia are the initial microcirculatory lesions in acute pancreatitis induced by sodium taurocholate leading to haemorrhagic necrosis. The non-specific detergent effect of sodium taurocholate and bile acids in general seems to be responsible for the initial changes due to the direct dissolution of cellular membranes.

### Changes of nutritive tissue perfusion

Acute pancreatitis is characterized by impairment of nutritive tissue perfusion as a consequence of gradually decreased capillary blood flow

and functional capillary density<sup>[60]</sup>. Reduction of capillary infusion volume and of functional capillary density has been observed with intravital microscopy and laser-Doppler flowmetry in the experiments of acute pancreatitis. In such experiments, capillaries are progressively excluded from perfusion starting 30min after the induction of pancreatitis, and with only few capillaries remaining perfused after 3h. At the same time, flow through the preferential pathways is maintained. Measurements of pancreatic blood flow during acute pancreatitis have ever yielded conflicting results. Some found no change or even increased blood flow, but most experiments have repeatedly demonstrated decreased total blood flow in acute pancreatitis. The perfusion values with an initial increase followed by a sharp decrease have been observed. Increased pancreatic blood flow is considered as a consequence of vasodilatation in acute inflammation. Because of the tremendous distributional disturbances of the microcirculation in the pancreas, however, measurements of total blood flow of the pancreatitis do not reflect proportionately the pathological status of different local regional perfusion within the pancreas. The pathological states, both the hyperemia and ischaemia, can be found at the same time in the different regions within the pancreas, thus emphasizing the importance of capillary blood flow measurement for accurate evaluation of microcirculatory blood flow changes. In most of the studies the degree of pancreatic hypoperfusion was found to be disproportionately more severe than the decrease in cardiac output at comparable intervals. Moreover it has been shown that a decrease in pancreatic perfusion cannot be prevented by adequate fluid therapy using Ringer's solution even though cardiovascular parameters are stabilized at the baseline level, proposing a specific mechanism of local microcirculatory ischaemic impairment<sup>[61-63]</sup>.

### Impairment of ischaemia/reperfusion and leukocyte adherence

Ischaemia/reperfusion of the pancreas with impairment of the microcirculation has attracted attention both in experimental and clinical studies of acute pancreatitis<sup>[64-73]</sup>. Ischaemia/reperfusion leads to the adherence of leukocytes to the vascular endothelium. In parallel with reduction of functional capillary density, an increase of heterogeneity of capillary perfusion has been noted. Primary capillary perfusion failure after onset of reperfusion is a characteristic microcirculatory feature of ischaemia and is called no-reflow phenomenon. Among various stimuli promoting leukocyte-endothelium interaction are ischaemia/reperfusion and formation of oxygen free radicals leading to rolling and adherence of leukocytes, the latter provoking the "reflow/paradox" phenomenon with loss of endothelial integrity and macromolecular leakage as an end result. Enhanced generation of oxygen radicals elicits ischaemia/reperfusion-induced leukocyte infiltration in the tissue, which is instrumental in the progression of acute pancreatitis. Degree of endothelial cell dysfunction and severity of leukocyte adherence is dependent upon the duration of ischaemia and reperfusion. Complete ischaemia/reperfusion of the pancreas induces extensive capillary stasis, i. e. pancreatic microcirculatory failure.

### Effect of hemorrhheological changes

Since blood viscosity is the inherent resistance of blood to flow, it is probable that the hemorrhheological changes might be important to acute necrotizing pancreatitis<sup>[74-84]</sup>. 188 Wistar rats were studied by measuring hemorrhheological and stereological parameters of pancreatic microvasculature. The results showed that increased blood viscosity, causing red blood cell aggregation with rouleaux formation, and decreased erythrocyte deformability are responsible for pancreatic microcirculatory disturbances and play an important role in the transition of oedematous pancreatitis to necrosis.

It has been noticed that the time points in the course of

experimental acute pancreatitis are extremely variable. This can be explained as investigators with various pancreatitis models, different infused substance, concentration, volume as well as intraductal pressure, the latter may be more important than the others. The high intraductal injection pressure results in an increased leakage of bile and a more generalized distribution in the interstitial space, even immediate hemorrhagic pancreatic necrosis, thus emphasizing the pathophysiological significance of experimental models in acute pancreatitis. In the low-pressure ductal perfusion model the etiological factor and the pathophysiological course are similar to those associated with the disease clinically.

## VASOACTIVE MEDIATORS IN ACUTE PANCREATITIS

### Bradykinin

Bradykinin probably exerts its influences upon microvessels via several pathways involving endothelial cells, including stimulating the formation and release of NO, arachidonic acid metabolites and tachykinins. Microcirculatory responses to bradykinin are biphasic: at low concentrations it causes vasodilatation, while at higher concentrations it causes vasoconstriction.

The role which bradykinin plays in microcirculatory impairment of acute pancreatitis is controversial. It was noticed that in sodium taurocholate-induced pancreatitis, the number of perfused capillaries was increased and capillary flow preserved and the mean venular leukocyte adherence decreased and histopathological change improved in icatibant (a B2 receptor antagonist)-treated rats; kinase II inhibitor captopril or exogenous bradykinin in addition to an otherwise effective dosage of icatibant resulted in microcirculatory stasis, extensive venular leukocyte adherence and severe histological damage, indicating that bradykinin may aggravate the microcirculatory disturbance<sup>[85,86]</sup>. But another study showed that B2 receptor antagonist increased the severity of acute pancreatitis, while lys-bradykinin substituting bradykinin didn't<sup>[87]</sup>.

### Platelet-activating factor(PAF)

PAF acts on microvascular diameter, permeability and leukocyte rolling, adhesion and migration through different mechanisms, including synthesis and release of NO and arachidonic acid metabolites, and upregulated expressions of ICAM-1 and CD11/CD18. Actions of PAF on microvasculature have the following features: constriction response of venules to PAF is stronger than that of the arterioles; its action on arteriolar diameter is biphasic.

It was observed that treatment with PAF receptor antagonist improved pancreatic capillary blood flow, reduced the severity of pancreatitis-associated endothelial barrier compromise and pancreatic leukocyte recruitment, suggesting that PAF is proinflammatory in pancreatitis<sup>[88-92]</sup>.

### Endothelin(ET)

There are three kinds of endothelins, and endothelin-1 is predominantly expressed by vascular endothelial cells. There are three types of endothelin receptors, and ETA receptor is endothelin-1 selective, and found mainly on vascular smooth muscle cells, mediating vasoconstriction; ETB is nonselective and expressed by endothelial cells; it mediates vasodilatation through the release of NO and prostacyclin. The action of endothelin-1 on microvessels is biphasic: at low concentration, it causes vasodilatation; at higher concentration, it causes sustained vasoconstriction.

Several experiments demonstrated that endothelin-1 was involved in the microcirculatory disturbance and in the development and progression of acute pancreatitis<sup>[93-99]</sup>. Administration of endothelin-1 after the caerulein injection decreased pancreatic blood flow significantly, aggravating microcirculatory disturbance. Topically

superfused endothelin-1 induced pancreatic microvascular deterioration and acinar cell injury similar to that induced by intraductal infusion of sodium taurocholate in rats<sup>[100]</sup>. Studies also showed that ETA receptor antagonist is protective in microcirculatory disturbance of acute pancreatitis<sup>[101]</sup>.

### Nitric oxide (NO)

NO is formed from L-arginine by NO synthase(NOS). cNOS(constitutive form) catalyzes formation of NO of physiological level. Catalytic activity of iNOS(inducible form) is stronger and lasts longer than that of cNOS, and NO of higher than physiological level is produced by iNOS. NO dilates blood vessels, but at higher concentrations it is cytotoxic.

Pancreatic NO level in acute pancreatitis may be decreased<sup>[102]</sup> or significantly elevated in different experiments. Intravenous administration of L-arginine to rats with hemorrhagic pancreatitis improved pancreatic blood flow and a meliorated the severity of pancreatitis in a dose-dependent manner, while nitro-L-arginine infusion to the rats with edematous pancreatitis caused a decrease in pancreatic blood flow and exacerbated pancreatitis, indicating that NO is protective<sup>[103-106]</sup>. However, some experiments showed that NO was not involved in the progression from edematous to hemorrhagic pancreatitis. Even microcirculatory changes were significantly alleviated in caerulein-induced pancreatitis pretreated with nitro-L-arginine, suggesting NO may be proinflammatory<sup>[107]</sup>; it was found that L-arginine improved the pancreatic microcirculation but worsened the microscopic alterations within the pancreas<sup>[108,109]</sup>.

### Adhesion molecules

Leukocyte-endothelial interaction is an important step in the development of acute pancreatitis. It was demonstrated by experiments that levels of ICAM-1, PECAM-1 and ELAM-1 were upregulated, and expressions of P- and E-selectin enhanced, and leukocytes became CD18-positive in acute pancreatitis<sup>[110,111]</sup>. Immunoneutralization of adhesion molecules was proven effective in the treatment of acute pancreatitis<sup>[112]</sup>. Administration of monoclonal antibody against ICAM-1 to rats with acute severe pancreatitis significantly enhanced capillary blood flow in the pancreas, reduced leukocyte rolling and stabilized capillary permeability<sup>[113]</sup>.

## PATHOGENESIS OF MICROCIRCULATORY FAILURE IN ACUTE PANCREATITIS

The pancreatic microcirculation is impaired in acute pancreatitis<sup>[114-124]</sup>. Capillary stasis may be due to a variety of mechanisms including hemoconcentration and intravascular coagulation, generation of oxygen free radicals in the microenvironment of the pancreatic ducto-acinar complex, increase in interstitial pressure, increase in leukocyte-endothelium interaction<sup>[125]</sup>, and local reduction of endothelial derived relaxation factor (nitrous oxide)<sup>[126]</sup>. An acinar abnormality may be the initiating factor arising from a combination of ductal obstruction and exocrine hypersecretion followed by an increase in intraductal pressure and leakage of enzymes into the pancreatic interstitium with release of zymogen and lysozymes.

Hemoconcentration and intravascular coagulation play an additional role in the development of pancreatic ischaemia in acute pancreatitis. The increased capillary permeability is the initial feature of experimental biliary pancreatitis, resulting in loss of fluid and cells into the pancreatic interstitium induced by osmolarity shifts either in the duct or extracellular fluid<sup>[127]</sup>. Local hemoconcentration takes place at the site of plasma sequestration, even if the systemic hematocrit is maintained at the initial level. In conjunction with the impairment of endothelium, intravascular coagulation occurs and causes a further decrease of blood fluidity. These changes are

aggravated by a systemic hypercoagulability in acute pancreatitis, probably due to thromboplastic material and activated trypsin gaining access to the systemic circulation.

The mechanism of oxygen free radical is important in acute pancreatitis of any causes<sup>[128]</sup> and is a direct sequel of biliopancreatic reflux at the onset of acute biliary pancreatitis. Besides disintegration of cell membranes by lipid peroxidation, free radicals trigger the extravasation of granulocytes into the surrounding parenchyma representing an early lesion in acute experimental pancreatitis. The initial margination of granulocytes in the capillaries may be a contributing factor in endothelial injury and impairment of capillary perfusion. Oxygen radicals mediate depletion of pancreatic sulphhydryl compounds with changes in both lipid peroxide and oxygen radical scavengers. Serum concentrations of vitamin C, a potent antioxidant, are depleted in acute pancreatitis so that synthetic ascorbic acid derivatives have been used as a free radical scavenger.

Postischemic intensive adherence of leukocytes to the endothelium of the venules and adhesive leukocytes forming plaques partially occluding the lumen of the venules have been observed within the reperfusional period in the experimental acute pancreatitis. This adhesive interaction is largely confined to postcapillary venules. And it is determined by a variety of factors such as expression of adhesion molecules on leukocytes and/or endothelial cells, products of leukocyte (superoxide) and endothelial cell (nitric oxide) activation and physical forces generated by the movement of blood along the vessel wall<sup>[129,130]</sup>. The firm adhesion of leukocytes that take place within postcapillary venules may increase the postcapillary pressure more than 200 folds, cause the passive dilatation of the capillaries and microcirculatory stasis. Many studies show that some compounds appear to be effective in reducing or abolishing leukocyte-endothelial cell adhesion, whereas some classical anti-inflammatory drugs such as indomethacin and aspirin actually promote leukocyte adhesion in the venules.

There may also be relation to lymphatic drainage<sup>[131]</sup>. Increase in local interstitial pressure as a consequence of obstructed lymph drainage further interferes with pancreatic microperfusion due to the venous outflow impairment. In the early period of acute experimental pancreatitis, dilated lymphatic vessels are visible macroscopically, and further progress of oedema with consequent focal hemorrhagic pancreatic necrosis is possible in case of insufficient lymphatic drainage. Experiment demonstrates that an increase of thoracic duct lymph flow followed by a pronounced and prolonged reduction. Erythrocytes originating from pancreatic interstitial hemorrhages were shown to enter and obstruct the microlymphatics.

## CONCLUSIONS

Recent advances in experimental research have helped witness the pathophysiology of acute pancreatitis. The phenomena of microcirculatory changes observed in acute experimental pancreatitis during the past few years gradually underlie the disturbance of the local microcirculation in acute pancreatitis, but several challenges remain. Still some questions remain unexplained concerning the mechanisms: (1) Which is the first event in the pathogenesis of acute pancreatitis? (2) Which factor determines the edematous or hemorrhagic necrotizing pancreatitis in a given experimental or clinical situation? (3) What is the role of impaired distribution of blood supply in early steps of acute pancreatitis? The potential vasoactive mediators responsible for the progression of the disease severity have largely remained subjecting to speculation and debate.

## REFERENCES

- Slavin J, Ghaneh P, Sutton R, Hartley M, Rowlands P, Garvey C, Hughes M, Neoptolemos J. Management of necrotizing pancreatitis. *World J Gastroenterol* 2001;7:476-481
- Wu K, Wang BX, Wang XP. Effects of clostridium butyricum on bacterial translocation in rats with acute necrotizing pancreatitis. *Shijie Huaren Xiaohua Zazhi* 2000;8:883-886
- Tu WF, Li JS, Zhu WM, Li ZD, Liu FN, Chen YM, Xu JG, Shao HF, Xiao GX, Li A. Influence of Glutamine and caecostomy/colonic irrigation on gut bacteria/endotoxin translocation in acute severe pancreatitis in pigs. *Shijie Huaren Xiaohua Zazhi* 1999;7:135-138
- Wu CT, Li ZL, Huang XC, Zhang ZL. Effect of Chinese medicine "Qing Yi Tang" and bifidobacterium mixture on intestinal bacterial translocation following acute necrotizing pancreatitis. *Shijie Huaren Xiaohua Zazhi* 1999;7:525-528
- Banks PA. Acute pancreatitis: medical and surgical management. *Am J Gastroenterol* 1994;89(8 Suppl):S78-85
- Kelly DM, McEntee GP, Delaney C, McGeeney KF, Fitzpatrick JM. Temporal relationship of acinar and microvascular changes in caerulein-induced pancreatitis. *Br J Surg* 1993;80:1174-1176
- Bockman DE. Microvasculature of the pancreas. Relation to pancreatitis. *Int J Pancreatol* 1992;12:11-21
- Klar E. Etiology and pathogenesis of acute pancreatitis. *Helv Chir Acta* 1992;59:7-16
- Waldner H. Vascular mechanisms to induce acute pancreatitis. *Eur Surg Res* 1992;24 Suppl 1:62-67
- Wu GD, Wu CW, Xu HB. Qingyitang decoction in treatment of 42 patients with severe acute pancreatitis. *Huaren Xiaohua Zazhi* 1998;6:619-621
- Sweiry JH, Mann GE. Pancreatic microvascular permeability in caerulein-induced acute pancreatitis. *Am J Physiol* 1991;261(4 Pt 1):G685-692
- Kusterer K, Enghofer M, Zendler S, Blochle C, Usadel KH. Microcirculatory changes in sodium taurocholate-induced pancreatitis in rats. *Am J Physiol* 1991;260(2 Pt 1):G346-351
- Klar E, Endrich B, Messmer K. Microcirculation of the pancreas. A quantitative study of physiology and changes in pancreatitis. *Int J Microcirc Clin Exp* 1990;9:85-101
- Wu CT, Li ZL. Anatomy of main pancreatic duct of hybrid dog and acute pancreatitis model. *Shijie Huaren Xiaohua Zazhi* 1999;7:62-63
- Chen HM, Shyr MH, Chen MF. Gabexate mesilate improves pancreatic microcirculation and reduces lung edema in a rat model of acute pancreatitis. *J For mos Med Assoc* 1997;96:704-709
- Gong ZH, Yuan YZ, Lou KX, Tu SP, Zhai ZK, Xu JY. Effects and mechanisms of somatostatin analogues on apoptosis of pancreatic acinar cells in acute pancreatitis in mice. *Shijie Huaren Xiaohua Zazhi* 1999;7:964-966
- Li TZ, Sun SQ, Sun DL. Serological studies on the metabolism of intercellular matrix in human acute pancreatitis. *Huaren Xiaohua Zazhi* 1998;6:1082-1083
- Chen HM, Hwang TL, Chen MF. The effect of gabexate mesilate on pancreatic and hepatic microcirculation in acute experimental pancreatitis in rats. *J Surg Res* 1996;66:147-53
- Huch K, Schmidt J, Schrott W, Sinn HP, Buhr H, Herfarth C, Klar E. Hyperoncotic dextran and systemic aprotinin in necrotizing rodent pancreatitis. *Scand J Gastroenterol* 1995;30:812-816
- Nishiwaki H, Satake K, Hiura A, Umeyama K. Effects of a newly synthesized pancreatic protease inhibitor (PATM) on pancreatic microcirculation in experimental acute pancreatitis. *Gastroenterol Jpn* 1989;24:177-180
- Li ZS, Xu GM, Sun ZX, Jin ZD, Zhou XP, Xie SQ, Li P. Early ERCP and endoscopic treatment in 66 patients with acute pancreatitis. *Huaren Xiaohua Zazhi* 1998;6:150-152
- Li ZS, Qian XD, Xu GM, Sun ZX, Zou XP, Xie SQ. Preventive effect of octreotide on hyperamylasemia and pancreatitis after ERCP. *Huaren Xiaohua Zazhi* 1998;6:617-618
- Kurzanov AN, Titova GP, Vinogradov VA, Aleinik VA, Gerasimov NF. Morphofunctional changes in the pancreas in response to dalargin under normal conditions and in experimental pancreatitis. *Biull Eksp Biol Med* 1988;105:445-447
- Lehtola A, Talja M, Puolakkainen P, Nordling S, Schroder T. Peritoneal lavage combined with volume therapy in porcine hemorrhagic pancreatitis. Effects on hemodynamics, microcirculation, and peritoneal morphology. *Scand J Gastroenterol* 1987;22:559-567
- Yuan YZ, Lou KX, Gong ZH, Tu SP, Zhai ZK, Xu JY. Effects and mechanisms of emodin on pancreatic tissue EGF expression in acute pancreatitis in rats. *Shijie Huaren Xiaohua Zazhi* 2001;9:127-130
- Xia SH, Zhao XY, Guo P, Da SP. Hemocirculatory disorder in dogs with severe acute pancreatitis and intervention of platelet activating factor antagonist. *Shijie Huaren Xiaohua Zazhi* 2001;9:550-554
- Wu CT, Li ZL. Effect of DAO on intestinal damage in acute necrotizing pancreatitis in dogs. *Shijie Huaren Xiaohua Zazhi* 1999;7:64-65

- 28 Zhao HP, Wang WX, Yang CW, Shou NY. Therapeutic effects of naltrexone in plasma endotoxin in experimental acute hemorrhagic necrotizing pancreatitis of rats. *Shijie Huaren Xiaohua Zazhi* 1999;7:400-402
- 29 Qin RY, Zou SQ, Wu ZD, Qiu FZ. Effect of splanchnic vascular perfusion on production of TNF $\alpha$  and OFR in rats with acute hemorrhagic necrotizing pancreatitis. *Huaren Xiaohua Zazhi* 1998; 6:831-833
- 30 Hirano T, Hirano K. Thromboxane A2 receptor antagonist prevents pancreatic microvascular leakage in rats with caerulein-induced acute pancreatitis. *Int J Surg Investig* 1999;1:203-210
- 31 Bhatia M, Saluja AK, Singh VP, Frossard JL, Lee HS, Bhagat L, Gerard C, Steer ML. Complement factor C5a exerts an anti-inflammatory effect in acute pancreatitis and associated lung injury. *Am J Physiol Gastrointest Liver Physiol* 2001;280:G974-978
- 32 Bhatia M, Brady M, Zagorski J, Christmas SE, Campbell F, Neoptolemos JP, Slavin J. Treatment with neutralising antibody against cytokine-induced neutrophil chemoattractant (CINC) protects rats against acute pancreatitis associated lung injury. *Gut* 2000; 47:838-844
- 33 Leung PS, Chan WP, Nobiling R. Regulated expression of pancreatic renin-angiotensin system in experimental pancreatitis. *Mol Cell Endocrinol* 2000;166:121-128
- 34 Gomez-Cambronero L, Camps B, de La Asuncion JG, Cerda M, Pellin A, Pallardo FV, Calvete J, Sweiry JH, Mann GE, Vina J, Sastre J. Pentoxifyline ameliorates cerulein-induced pancreatitis in rats: role of glutathione and nitric oxide. *J Pharmacol Exp Ther* 2000;293: 670-676
- 35 Hirota M, Ogawa M. Shock and its mediators. *Nippon Geka Gakkai Zasshi* 1999;100:667-673
- 36 al-Eryani S, Payer J, Huorka M, Duris I. Etiology and pathogenesis of acute pancreatitis. *Bratisl Lek Listy* 1998;99:303-311
- 37 Plusczyk T, Westermann S, Rathgeb D, Feifel G. Acute pancreatitis in rats: effects of sodium taurocholate, CCK-8, and Sec on pancreatic microcirculation. *Am J Physiol* 1997;272(2 Pt 1):G310-320
- 38 Sakai Y, Hayakawa T, Kondo T, Shibata T, Kitagawa M, Sobajima H, Naruse S, Ohnishi ST. Protective effects of a prostaglandin E1 oligomer on taurocholate-induced rat pancreatitis. *J Gastroenterol Hepatol* 1992;7:591-595
- 39 Vollmar B, Waldner H, Schmand J, Conzen PF, Goetz AE, Habazettl H, Schweiberer L, Brendel W. Oleic acid induced pancreatitis in pigs. *J Surg Res* 1991;50:196-204
- 40 Lehtola A, Kivilaakso E, Puolakkainen P, Karonen SL, Lempinen M, Schroder T. Effects of dextran 70 versus crystalloids in the microcirculation of porcine hemorrhagic pancreatitis. *Surg Gynecol Obstet* 1986; 162:556-562
- 41 Kaplan MH. Pathogenesis of pancreatitis: a unified concept. *Int J Pancreatol* 1986;1:5-8
- 42 Sunamura M, Yamauchi J, Shibuya K, Chen HM, Ding L, Takeda K, Kobari M, Matsuno S. Pancreatic microcirculation in acute pancreatitis. *J Hepatobil Pancreat Surg* 1998;5:62-68
- 43 Plusczyk T, Rathgeb D, Westermann S, Feifel G. Somatostatin attenuates microcirculatory impairment in acute sodium taurocholate-induced pancreatitis. *Dig Dis Sci* 1998;43:575-585
- 44 Schmidt J, Klar E. Etiology and pathophysiology of acute pancreatitis. *Ther Umsch* 1996;53:322-332
- 45 Hoffmann TF, Leiderer R, Waldner H, Arbogast S, Messmer K. Ischemia reperfusion of the pancreas: a new *in vivo* model for acute pancreatitis in rats. *Res Exp Med (Berl)* 1995; 195:125-144
- 46 Sendur R, Pawlik WW. Vascular factors in the mechanism of acute pancreatitis. *Przegl Lek* 1996;53:41-45
- 47 Moolenaar W, Lamers CB. Cholesterol crystal embolization and the digestive system. *Scand J Gastroenterol Suppl* 1991;188:69-72
- 48 Hegewald G, Nikulin A, Gmaz-Nikulin E, Plamenac P, Barenwald G. Ultrastructural changes of the human pancreas in acute shock. *Pathol Res Pract* 1985;179:610-615
- 49 Klar E, Schrott W, Foitzik T, Buhr H, Herfarth C, Messmer K. Impact of microcirculatory flow pattern changes on the development of acute edematous and necrotizing pancreatitis in rabbit pancreas. *Dig Dis Sci* 1994;39:2639-2644
- 50 Zhou ZG, Gao XH. Morphology of pancreatic microcirculation in the monkey: light and scanning electron microscopic study. *Clin Anat* 1995; 8:190-201
- 51 Zhou Z, Zeng Y, Yang P, Cheng Z, Zhao J, Shu Y, Gao X, Yan L, Zhang Z. Structure and function of pancreatic microcirculation. *Shengwu Yiyue Gongchengxue Zazhi* 2001;18:195-200
- 52 Furukawa M, Kimura T, Sumii T, Yamaguchi H, Nawata H. Role of local pancreatic blood flow in development of hemorrhagic pancreatitis induced by stress in rats. *Pancreas* 1993;8:499-505
- 53 Plusczyk T, Westermann S, Bersal B, Menger M, Feifel G. Temporary pancreatic duct occlusion by ethibloc: cause of microcirculatory shutdown, acute inflammation, and pancreas necrosis. *World J Surg* 2001;25:432-437
- 54 Bassi D, Kollias N, Fernandez-del Castillo C, Foitzik T, Warshaw AL, Rattner DW. Impairment of pancreatic microcirculation correlates with the severity of acute experimental pancreatitis. *J Am Coll Surg* 1994;179:257-263
- 55 Banerjee AK, Galloway SW, Kingsnorth AN. Experimental models of acute pancreatitis. *B J Surg* 1994; 81:1096-1103
- 56 Klar E, Werner J. New pathophysiologic knowledge about acute pancreatitis. *Chirurg* 2000;71:253-264
- 57 Onizuka S, Ito M, Sekine I, Tsunoda T, Eto T. Spontaneous pancreatitis in spontaneously hypertensive rats. *Pancreas* 1994;9:54-61
- 58 Skoromnyi AN, Starosek VN. Hemodynamic changes in the liver, kidney, small intestine and pancreas in experimental acute pancreatitis. *Klin Khir* 1998;12:46-48
- 59 Kusterer K, Poschmann T, Friedemann A, Enghofer M, Zendler S, Usadel KH. Arterial constriction, ischemia-reperfusion, and leukocyte adherence in acute pancreatitis. *Am J Physiol* 1993; 265(1 Pt 1): G165-171
- 60 Kerner T, Vollmar B, Menger MD, Waldner H, Messmer K. Determinants of pancreatic microcirculation in acute pancreatitis in rats. *J Surg Res* 1996;62:165-171
- 61 Chen HM, Shyr MH, Ueng SW, Chen MF. Hyperbaric oxygen therapy attenuates pancreatic microcirculatory derangement and lung edema in an acute experimental pancreatitis model in rats. *Pancreas* 1998;17: 44-49
- 62 Klar E, Messmer K, Warshaw AL, Herfarth C. Pancreatic ischaemia in experimental acute pancreatitis: mechanism, significance and therapy. *Br J Surg* 1990;77:1205-1210
- 63 Knol JA, Inman MG, Strodel WE, Eckhauser FE. Pancreatic response to crystalloid resuscitation in experimental pancreatitis. *J Surg Res* 1987;43:387-392
- 64 Obermaier R, Benz S, Kortmann B, Benthues A, Ansorge N, Hopt UT. Ischemia/reperfusion-induced pancreatitis in rats: a new model of complete normothermic *in situ* ischemia of a pancreatic tail-segment. *Clin Exp Med* 2001;1:51-59
- 65 Benz S, Bergt S, Obermaier R, Wiessner R, Pfeffer F, Schareck W, Hopt UT. Impairment of microcirculation in the early reperfusion period predicts the degree of graft pancreatitis in clinical pancreas transplantation. *Transplantation* 2001; 27:71:759-763
- 66 Mayer H, Schmidt J, Thies J, Ryschich E, Gebhard MM, Herfarth C, Klar E. Characterization and reduction of ischemia/reperfusion injury after experimental pancreas transplantation. *J Gastrointest Surg* 1999;3:162-166
- 67 von Dobschuetz E, Hoffmann T, Messmer K. Inhibition of neutrophil proteinases by recombinant serpin Lex032 reduces capillary no-reflow in ischemia/reperfusion-induced acute pancreatitis. *J Pharmacol Exp Ther* 1999;290:782-788
- 68 von Dobschuetz E, Hoffmann T, Engelschalk C, Messmer K. Effect of diaspirin cross-linked hemoglobin on normal and postischemic microcirculation of the rat pancreas. *Am J Physiol* 1999;276(6 Pt 1): G1507-1514
- 69 Vollmar B, Janata J, Yamauchi J, Wolf B, Heuser M, Menger MD. Exocrine, but not endocrine, tissue is susceptible to microvascular ischemia/reperfusion injury following pancreas transplantation in the rat. *Transpl Int* 1999;12:50-55
- 70 Benz S, Pfeffer F, Adam U, Schareck W, Hopt UT. Impairment of pancreatic microcirculation in the early reperfusion period during simultaneous pancreas-kidney transplantation. *Transpl Int* 1998;11 Suppl 1:S433-435
- 71 Benz S, Schnabel R, Morgenroth K, Weber H, Pfeffer F, Hopt UT. Ischemia/reperfusion injury of the pancreas: a new animal model. *J Surg Res* 1998;75:109-115
- 72 Hoffmann TF, Leiderer R, Harris AG, Messmer K. Ischemia and reperfusion in pancreas. *Microsc Res Tech* 1997;37:557-571
- 73 Menger MD, Bonkhoff H, Vollmar B. Ischemia-reperfusion-induced pancreatic microvascular injury. An intravital fluorescence microscopic study in rats. *Dig Dis Sci* 1996;41:823-830
- 74 Schmidt J, Huch K, Mithofer K, Hotz HG, Sinn HP, Buhr HJ, Warshaw AL, Herfarth C, Klar E. Benefits of various dextrans after delayed therapy in necrotizing pancreatitis of the rat. *Intensive Care Med* 1996; 22:1207-1213
- 75 Hotz HG, Schmidt J, Ryschich EW, Foitzik T, Buhr HJ, Warshaw AL, Herfarth C, Klar E. Isovolemic hemodilution with dextran prevents contrast medium induced impairment of pancreatic microcirculation in necrotizing pancreatitis of the rat. *Am J Surg* 1995;169:161-166

- 76 Klar E, Foitzik T, Buhr H, Messmer K, Herfarth C. Isovolemic hemodilution with dextran 60 as treatment of pancreatic ischemia in acute pancreatitis. Clinical practicability of an experimental concept. *Ann Surg* 1993;217:369-374
- 77 Yan LN, Lei ZM, Cui XZ, Chen HQ, Yang YT, Li L, Tan JS, Chen LL, Wu HB, Li KL. The role of hemorheologic disturbance in experimental acute pancreatitis. *Huaxi Yike Daxue Xuebao* 1993;24:71-74
- 78 Klar E, Mall G, Messmer K, Herfarth C, Rattner DW, Warshaw AL. Improvement of impaired pancreatic microcirculation by isovolemic hemodilution protects pancreatic morphology in acute biliary pancreatitis. *Surg Gynecol Obstet* 1993;176:144-150
- 79 Schmidt J, Fernandez-del Castillo C, Rattner DW, Lewandrowski KB, Messmer K, Warshaw AL. Hyperoncotic ultrahigh molecular weight dextran solutions reduce trypsinogen activation, prevent acinar necrosis, and lower mortality in rodent pancreatitis. *Am J Surg* 1993;165:40-44
- 80 Kusterer K, Enghofer M, Poschmann T, Usadel KH. The effect of somatostatin, gabexate mesilate and dextran 40 on the microcirculation in sodium taurocholate-induced pancreatitis. *Acta Physiol Hung* 1992;80:407-415
- 81 Yan LN, Wei JJ, Wu HG, Chen HQ, Zhong GH, Li L, Chen LL, Li KL, Tan JS. The role of hemorheologic changes in the pathogenesis of acute hemorrhagic necrotizing pancreatitis. *Huaxi Yike Daxue Xuebao* 1990;21:25-29
- 82 Klar E, Herfarth C, Messmer K. Therapeutic effect of isovolemic hemodilution with dextran 60 on the impairment of pancreatic microcirculation in acute biliary pancreatitis. *Ann Surg* 1990;211:346-353
- 83 Aleksandrova NP, Petukhov EB, Riabova SS. Blood rheology and microcirculation in the dynamics of acute experimental pancreatitis. *Biull Eksp Biol Med* 1988;105:106-108
- 84 Becker H, Senninger N. Hemorrhagic pancreatitis: effect of dextran 40 and plasma on microcirculation disorders of the pancreas. *Langenbecks Arch Chir* 1985;365:57-67
- 85 Bloechle C, Kusterer K, Kuehn RM, Schneider C, Knoefel WT, Izbicki JR. Inhibition of bradykinin B2 receptor preserves microcirculation in experimental pancreatitis in rats. *Am J Physiol* 1998; 274(1 Pt 1): G42-51
- 86 Hoffmann T, Kubler J, Messmer K. Bradykinin antagonism in ischemia and reperfusion of the pancreas. *Zentralbl Chir* 1996;121:412-422
- 87 Weidenbach H, Lerch MM, Gress TM, Pfaff D, Turi S, Adler G. Vasoactive mediators and the progression from oedematous to necrotising experimental acute pancreatitis. *Gut* 1995;37:434-440
- 88 Flickinger BD, Olson MS. Localization of the platelet-activating factor receptor to rat pancreatic microvascular endothelial cells. *Am J Pathol* 1999;154:1353-1358
- 89 Foitzik T, Hotz HG, Eibl G, Hotz B, Kirchengast M, Buhr HJ. Therapy for microcirculatory disorders in severe acute pancreatitis: effectiveness of platelet-activating factor receptor blockade vs. endothelin receptor blockade. *J Gastrointest Surg* 1999;3:244-251
- 90 Wang X, Sun Z, Borjesson A, Haraldsen P, Aldman M, Deng X, Leveau P, Andersson R. Treatment with lexipafant ameliorates the severity of pancreatic microvascular endothelial barrier dysfunction in rats with acute hemorrhagic pancreatitis. *Int J Pancreatol* 1999;25: 45-52
- 91 Ji Z, Wang B, Li S. The role of platelet activating factor in mesenterioangial microcirculatory disturbance complicated with acute pancreatitis in rats. *Zhonghua Yixue Zazhi* 1995;75:139-140
- 92 Travis SP, Jewell DP. The role of platelet-activating factor in the pathogenesis of gastrointestinal disease. *Prostagl Leukot Essent Fatty Acids* 1994;50:105-113
- 93 Plusczyk T, Bersal B, Menger MD, Feifel G. Differential effects of ET-1, ET-2, and ET-3 on pancreatic microcirculation, tissue integrity, and inflammation. *Dig Dis Sci* 2001;46:1343-1351
- 94 Foitzik T, Faulhaber J, Hotz HG, Kirchengast M, Buhr HJ. Endothelin mediates local and systemic disease sequelae in severe experimental pancreatitis. *Pancreas* 2001;22:248-254
- 95 Foitzik T, Eibl G, Hotz HG, Faulhaber J, Kirchengast M, Buhr HJ. Endothelin receptor blockade in severe acute pancreatitis leads to systemic enhancement of microcirculation, stabilization of capillary permeability, and improved survival rates. *Surgery* 2000;128:399-407
- 96 Foitzik T, Eibl G, Buhr HJ. Therapy for microcirculatory disorders in severe acute pancreatitis: comparison of delayed therapy with ICAM-1 antibodies and a specific endothelin A receptor antagonist. *J Gastrointest Surg* 2000;4:240-246
- 97 Foitzik T, Hotz HG, Hot B, Kirchengast M, Buhr HJ. Endothelin-1 mediates the alcohol-induced reduction of pancreatic capillary blood flow. *J Gastrointest Surg* 1998;2:379-384
- 98 Foitzik T, Faulhaber J, Hotz HG, Kirchengast M, Buhr HJ. Endothelin receptor blockade improves fluid sequestration, pancreatic capillary blood flow, and survival in severe experimental pancreatitis. *Ann Surg* 1998;228:670-675
- 99 Liu XH, Kimura T, Ishikawa H, Yamaguchi H, Furukawa M, Nakano I, Kinjoh M, Nawata H. Effect of endothelin-1 on the development of hemorrhagic pancreatitis in rats. *Scand J Gastroenterol* 1995;30:276-282
- 100 Plusczyk T, Bersal B, Westermann S, Menger M, Feifel G. ET-1 induces pancreatitis-like microvascular deterioration and acinar cell injury. *J Surg Res* 1999; 85: 301-310
- 101 Liu X, Nakano I, Ito T, Kimura T, Nawata H. Is endothelin-1 an aggravating factor in the development of acute pancreatitis? *Chin Med J (Engl)* 1999;112:603-607
- 102 Shibuya K, Sunamura M, Yamauchi J, Takeda K, Kobari M, Matsuno S. Analysis of the derangement of the pancreatic microcirculation in a rat cerulein pancreatitis model using an intravital microscope system. *Tohoku J Exp Med* 1996; 180: 173-186
- 103 Vollmar B, Janata J, Yamauchi JJ, Menger MD. Attenuation of microvascular reperfusion injury in rat pancreas transplantation by L-arginine. *Transplantation* 1999;67:950-955
- 104 Dobosz M, Hac S, Mionskowska L, Dobrowolski S, Wajda Z. Microcirculatory disturbances of the pancreas in cerulein-induced acute pancreatitis in rats with reference to L-arginine, heparin, and procaine treatment. *Pharmacol Res* 1997;36:123-128
- 105 Dobosz M, Hac S, Wajda Z. Does nitric oxide protect from microcirculatory disturbances in experimental acute pancreatitis in rats? *Int J Microcirc Clin Exp* 1996;16:221-226
- 106 Liu X, Nakano I, Yamaguchi H, Ito T, Goto M, Koyanagi S, Kinjoh M, Nawata H. Protective effect of nitric oxide on development of acute pancreatitis in rats. *Dig Dis Sci* 1995;40:2162-2169
- 107 Dobosz M, Wajda Z, Hac S, Mysliwska J, Mionskowska L, Bryl E, Roszkiewicz A, Mysliwski A. Heparin and nitric oxide treatment in experimental acute pancreatitis in rats. *Forum (Genova)* 1998;8: 303-310
- 108 Dobosz M, Wajda Z, Hac S, Mysliwska J, Bryl E, Mionskowska L, Roszkiewicz A, Mysliwski A. Nitric oxide, heparin and procaine treatment in experimental cerulein-induced acute pancreatitis in rats. *Arch Immunol Ther Exp (Warsz)* 1999;47:155-160
- 109 Hac DS, Mionskowska L, Dobrowolski S, Dymecki D, Makarewicz W, Wajda Z. Microcirculation disorders of the pancreas in cerulein induced acute pancreatitis in rats with regard to nitrogen oxide and heparin. *Wiad Lek* 1997;50(Pt 2):108-114
- 110 Lundberg AH, Granger DN, Russell J, Sabek O, Henry J, Gaber L, Kotb M, Gaber AO. Quantitative measurement of P- and E-selectin adhesion molecules in acute pancreatitis: correlation with distant organ injury. *Ann Surg* 2000; 231: 213-222
- 111 Sunamura M, Shibuya K, Yamauchi J, Matsuno S. Microcirculatory derangement and ischemia of the pancreas. *Nippon Geka Gakkai Zasshi* 1999;100:342-346
- 112 Wang X, Sun Z, Borjesson A, Andersson R. Inhibition of platelet-activating factor, intercellular adhesion molecule 1 and platelet endothelial cell adhesion molecule 1 reduces experimental pancreatitis-associated gut endothelial barrier dysfunction. *Br J Surg* 1999;86: 411-416
- 113 Frossard JL, Saluja A, Bhagat L, Lee HS, Bhatia M, Hofbauer B, Steer ML. The role of intercellular adhesion molecule 1 and neutrophils in acute pancreatitis and pancreatitis-associated lung injury. *Gastroenterology* 1999;116:694-701
- 114 Kaska M, Pospisilova B, Slizova D. Pathomorphological changes in microcirculation of pancreas during experimental acute pancreatitis. *Hepatogastroenterology* 2000;47:1570-1574
- 115 Knoefel WT, Kollias N, Warshaw AL, Waldner H, Nishioka NS, Rattner DW. Pancreatic microcirculatory changes in experimental pancreatitis of graded severity in the rat. *Surgery* 1994;116: 904-913
- 116 Foitzik T, Bassi DG, Fernandez-del Castillo C, Warshaw AL, Rattner DW. Intravenous contrast medium impairs oxygenation of the pancreas in acute necrotizing pancreatitis in the rat. *Arch Surg* 1994;129: 706-711
- 117 Kelly DM, McEntee GP, McGeeney KF, Fitzpatrick JM. Microvasculature of the pancreas, liver, and kidney in cerulein-induced pancreatitis. *Arch Surg* 1993;128:293-295
- 118 Garcia-Cano J, Vazquez Rodriguez de Alba J, Garcia Cabezas J, Diaz-Rubio M. Acute pancreatitis in thrombotic thrombocytopenic purpura. Apropos 2 cases. *An Med Interna* 1992;9:551-553
- 119 Waldner H, Schmand J, Vollmar B, Goetz A, Conzen P, Schweiberer L, Brendel W. Pancreatic circulation in experimental biliary pancreatitis. *Langenbecks Arch Chir* 1990;375:112-118
- 120 Gress TM, Arnold R, Adler G. Structural alterations of pancreatic microvasculature in cerulein-induced pancreatitis in the rat. *Res Exp Med*

- (Berl) 1990;190:401-412
- 121 McEntee G, Leahy A, Cottell D, Dervan P, McGeeney K, Fitzpatrick JM. Three-dimensional morphological study of the pancreatic microvasculature in caerulein-induced experimental pancreatitis. *Br J Surg* 1989;76:853-855
- 122 Nishiwaki H, Satake K, Ko I, Umeyama K. Pancreatic microcirculation in acute pancreatitis of dogs. *Nippon Geka Gakkai Zasshi* 1988;89:238-244
- 123 Nuutinen P, Kivisaari L, Standertskjold-Nordenstam CG, Lempinen M, Schroder T. Microangiography of the pancreas in experimental hemorrhagic pancreatitis. *Eur J Radiol* 1986;6:187-190
- 124 Nuutinen P, Kivisaari L, Standertskjold-Nordenstam CG, Lempinen M, Schroder T. Microangiography of the pancreas in experimental oedemic and haemorrhagic pancreatitis. *Scand J Gastroenterol Suppl* 1986;126:12-17
- 125 Chen HM, Sunamura M, Shibuya K, Yamauchi JI, Sakai Y, Fukuyama S, Mikami Y, Takeda K, Matsuno S. Early microcirculatory derangement in mild and severe pancreatitis models in mice. *Surg Today* 2001;31:634-642
- 126 Menger MD, Plusczyk T, Vollmar B. Microcirculatory derangements in acute pancreatitis. *J Hepatobiliary Pancreat Surg* 2001;8:187-194
- 127 Andre EA, Costa PL, Guarita DR, Meirelles Filho JS, Laudanna AA. Changes in capillary permeability during severe experimental acute pancreatitis in rats. *Rev Hosp Clin Fac Med Sao Paulo* 1996;51:184-188
- 128 Petersson U, Kallen R, Montgomery A, Borgstrom A. Role of oxygen-derived free radicals in protease activation after pancreas transplantation in the pig. *Transplantation* 1998;65:421-426
- 129 Inagaki H, Nakao A, Kurokawa T, Nonami T, Harada A, Takagi H. Neutrophil behavior in pancreas and liver and the role of nitric oxide in rat acute pancreatitis. *Pancreas* 1997;15:304-309
- 130 Toyama MT, Lewis MP, Kusske AM, Reber PU, Ashley SW, Reber HA. Ischaemia-reperfusion mechanisms in acute pancreatitis. *Scand J Gastroenterol Suppl* 1996;219:20-23
- 131 Kul'chitskii KI, Zurnadzhi IN, Blagodarov VN, Bogdanova GI, Shchitov VS, Ponomareva VP. The lymph and blood microcirculatory beds of the pancreas in the early period of experimental acute pancreatitis. *Lik Sprava* 1994;87-89

Edited by Wu XN